

# Ocean Colour Remote Sensing in Turbid Waters

## Lecture 2: Introduction to computer exercise “The Colour of Water”

by Kevin Ruddick

# Overview of this lecture

- Objective: introduce the HYPERTEACH ocean colour model as basis for exercise session
- NB. This is an approximate model for educational purposes only
- NOT for ocean colour data processing
- NOT for research grade publications
- JUST for understanding first order variability of marine reflectance
- CONDITIONS of USE:
  - I will not hold anyone responsible for mis-use, etc.
  - I will not use this for ocean colour data processing or research grade publications - for that I will use accurate radiative transfer models such as HYDROLIGHT (water) or 6SV (atmosphere)
  - I will use this model for quickly understanding ocean colour variability
  - I will not cheat and go straight to the answers
  - I will think of ways this could be improved for educational purposes and help by providing suggestions

We Accept

# Variation of reflectance with IOPs

- Gordon-Morel type approximate reflectance model

$$R_{rs}(\lambda) = \gamma' \frac{b_b(\lambda)}{a(\lambda) + b_b(\lambda)}$$

- For all but most reflective water, relation is **linear**:

$$b_b \ll a \quad \Rightarrow \quad R_{rs}(\lambda) = \gamma' \frac{b_b(\lambda)}{a(\lambda)}$$

- (NB This model is not appropriate for high reflectance)

# Decomposition of IOPs: absorption

- The total absorption can be decomposed into a linear sum of (mutually exclusive) components:

$$a(\lambda) = a_w(\lambda) + a_\phi(\lambda) + \overbrace{a_{CDOM}(\lambda) + a_{NAP}(\lambda)}^{(total) \text{ yellow substance } a_Y(\lambda)}$$

Pure water

Phytoplankton

Coloured Dissolved Organic Matter

Non-algae particles




# Optical properties of pure sea water (1/3)

- Backscatter of pure sea water (includes bubbles?):
  - Generally low, especially for green-red


$$b_{bw} = 0.5 * 0.00288 * \left( \frac{\lambda}{500nm} \right)^{-4.32} \quad [\text{Morel, 1974}]$$

Pure water backscatter



# Optical properties of pure sea water (2/3)

- Absorption of pure sea water:
  - Dominant absorber (except in very turbid waters) for red and especially near infrared




# Optical properties of pure water (3/3)

- If water contains no other constituents (no phytoplankton or other particles, no coloured dissolved organic matter) then:

$$R_{rs}(\lambda) = \gamma' \frac{b_b(\lambda)}{a(\lambda)} \approx 0.069 \frac{b_{bw}(\lambda)}{a_w(\lambda)}$$

Pure water reflectance



- Not a realistic case, but useful extreme case (blue/violet water)



# Optical properties of phytoplankton (1/2)

- Backscatter of phytoplankton:

- Main backscatterer in open ocean, relatively flat spectrum


$$\nu \in (-0.65, 0)$$

$$b_{b\phi} = \left\{ 0.002 + 0.01 * [0.50 - 0.25 \log_{10} C] \left( \frac{\lambda}{550nm} \right)^{\nu} \right\} * 0.416 * C^{0.766}$$

[Morel and Maritorena, 2001]


Phytoplankton backscatter

$C = \text{Chl a}$



# Optical properties of phytoplankton (2/2)


- Absorption of phytoplankton:
  - Main absorber in open ocean, spectral features in blue and red
  - Phyto absorption proportional to Chl a (first approximation)
  - Tabulated spectra given as function of Chl a [Bricaud et al, 1995]



# Coloured Dissolved Organic Matter (CDOM)

- CDOM=humic/fulvic acids from degradation of **terrestrial** or **marine** vegetation (correlated with **salinity** or **phytoplankton**)
  - neg. backscatter, absorbs strongly in blue: « yellow » substance
  - can be main absorber in coastal waters with high river input but low suspended matter e.g. parts of Baltic Sea, Black Sea


Coloured Dissolved Organic Matter (CDOM) absorption



# Optical properties of non-algal particles (1/2)


- Non-algal particles (NAP) may have diverse nature and origin: e.g. mineral particles (coastal/bottom erosion, river outflow), detrital particles (decayed phytoplankton)
- Backscatter relatively flat spectrally,  $\propto$  NAP concentration, can be main backscatterer in coastal and estuarine waters

Non-algae particle backscatter




# Optical properties of non-algal particles (2/2)


- Absorption of non-algal particles is strong in blue (like CDOM) with exponential decrease to higher wavelengths: « particulate » yellow substance
- Proportional to conc. of non-algae particles



# From water constituents to reflectance via IOPs



# Example reflectance spectra



# Exceptions

- Assumes:
  - No bottom reflectance
  - No inelastic scattering (fluorescence, Raman, bioluminescence)
  - Vertically homogeneous (no stratification, no deep CHL max, etc.)



# Make your own reflectance spectra

- Now follow the exercises and make your own reflectance spectra ...